arXiv:1605.04156v1 [cs.CG] 13 May 2016

Rational Bézier Curves Approximated by Bernstein-Jacobi Hybrid Polynomial Curves

Mao Shi[a]

[a]*School of Mathematics and Information Science of Shaanxi Normal University, Xian 710062, China*

Abstract

In this paper, we propose a linear method for $C^{(r,s)}$ approximation of rational Bézier curve with arbitrary degree polynomial curve. Based on weighted least-squares, the problem be converted to an approximation between two polynomial curves. Then applying Bernstein-Jacobi hybrid polynomials, we obtain the resulting curve. In order to reduce error, degree reduction method for Bézier curve is used. A error bound between rational Bézier curve and Bézier curve is presented. Finally, some examples and figures were offered to demonstrate the efficiency, simplicity, and stability of our methods.

Keywords: Rational Bézier curves; Jacobi polynomials; Polynomial approximation; Weighted least-squares method; Degree reduction method

1. Introduction

Rational Bézier curve is an important tools for Computer-Aided Design (CAD) to present conic precisely, but its calculus is complex or may not exist for high degree forms, so many CAD systems only use polynomial expressions to deal with parametric curves [1]. Therefore, it is certainly worthy researching how to approximate rational Bézier curves by polynomial parametric curves.

*Email: shimao@snnu.edu.cn

Many authors have studied the field since 1991. Sederberg and Kakimoto [2] first presented the hybrid algorithm for polynomial approximation of rational curve. Hereafter, Wang and Sederberg[3] analyzed its convergence condition and put forward a recursive algorithm. Lee and Park [4] converted the problem to control points approximation of two Bézier curves. Using dual constrained Bernstein polynomials and Chebyshev polynomials, Lewanowicz etc.[5] derived a polynomial curve approximating a rational Bézier curve with endpoints constraints in the L_2-norm and exploited their recursive properties. Shi and Deng [6] introduced weighted least squares approximation to the field. Hu and Xu [7] proposed a reparameterization-based method for polynomial approximating rational Bézier curves.

The degree reduction of Bézier curve is of particular importance in the field of CAD [8]. It was mainly used in data exchange, data compression and data comparsion [9]. Jacobi polynomials are a key tool for degree reduction of Bézier curve, which have been studied by many researchers [10, 11, 12, 13, 14, 15]. The main ideas of this method are first converting Bernstein polynomials to Jacobi Polynomials, then truncating the highest coefficient or using least squares to obtain the best approximation. Finally converting back the Jacobi series to Bernstein form for getting the solutions.

In this paper, we direclty apply Jacobi polynomials to achieve polynomial approximation of rational Bézier curves with $C^{(r,s)}$- continuity. To avoid calculus of rational Bézier curves, we convert the problem to an approximation problem of two polynomial curves by weighted least-squares, and we find that if using the degree reduction method, the error between rational Bézier curve and bézier curve sometimes can be reduced. Through numerical examples, we also show that our method is feasible and effective.

The paper is structured as follows. Section 2 presents some basic concepts and properties regarding the problem of the constrained Bézier approximation of a rational Bézier curve. Section 3 brings a complete solution to the

problem formulated in the L_2 norm. Section 4 presents some numerical examples to verify the accuracy and effectiveness of the method.

2. Preliminary

Given a rational Bézier curve of degree n

$$\mathbf{x(t)} = \frac{\mathbf{p}(t)}{\omega(t)} = \frac{\sum\limits_{i=0}^{n} \omega_i \mathbf{p}_i B_i^n(t)}{\sum\limits_{i=0}^{n} \omega_i B_i^n(t)} = \frac{\sum\limits_{i=0}^{n} \mathbf{P}_i B_i^n(t)}{\sum\limits_{i=0}^{n} \omega_i B_i^n(t)} \tag{1}$$

where $\mathbf{p}_i$ are called control points, $B_i^n(t)$ are Bernstein basis of degree n, $\omega_i > 0$ are weights and $\mathbf{P}_i = \omega_i \mathbf{p}_i$, the following curve

$$\tilde{\mathbf{y}}(t) = \sum_{i=0}^{r} \mathbf{q}_i B_i^m(t) + t^{r+1}(1-t)^{s+1} \sum_{j=0}^{m-(r+s+2)} \tilde{\mathbf{q}}_j J_j^{(r+1,s+1)}(2t-1) + \sum_{i=m-s}^{m} \mathbf{q}_i B_i^m(t), \tag{2}$$

is called a Bernstein-Jacobi hybrid curve approximation to the rational Bézier curve $\mathbf{x}(t)$ in L_2 norm, if they satisfy weighted least squares

$$min \quad d = \int_0^1 \rho(t) \left\| \mathbf{x}(t) - \tilde{\mathbf{y}}(t) \right\|^2 dt, \tag{3}$$

where $\rho(t) > 0$, $J_j^{(r+1,s+1)}(2t-1)$ are Jacobi polynomial, $\mathbf{q}_i$ are constrained control points and $\tilde{\mathbf{q}}_j$ are Jacobi control points.

For the derivation of our polynomial method, we need the following properties[9, 16, 17].

1. The definite integral of Bernstein basis is

$$\int_0^1 B_i^n(t) = \frac{1}{n+1}. \tag{4}$$

2. Let $M = m - (r + s + 2)$ and $k = 0, ..., M$. The weighted Jacobi polynomial can be represented in Bernstein forms as

$$t^{\alpha+1}(1-t)^{\beta+1}J_k^{(r+1,s+1)}(2t-1)$$
$$= \begin{cases} \sum_{i=0}^{k}(-1)^{k+i}\frac{\binom{k+r+1}{i}\binom{k+s+1}{k-i}}{\binom{k}{i}}B_i^k(t), & \alpha=\beta=-1, \\ \sum_{i=0}^{k}\sum_{j=i}^{M+i-k}(-1)^{k+i}\frac{\binom{k+r+1}{i}\binom{k+s+1}{k-i}\binom{M-k}{j-i}}{\binom{m}{r+j+1}}B_{r+j+1}^m(t), & \alpha=r>-1,\ \beta=s>-1, \end{cases} \tag{5}$$

3. Let $f(t) = \sum_{i=0}^{m} f_i B_i^m(t)$ and $g(t) = \sum_{j=0}^{n} g_j B_j^n(t)$ are the two Bernstein polynomial functions of degree m and n with coefficients f_i and g_j. Their product is a degree $m+n$ polynomial

$$f(t)g(t) = \sum_{i=0}^{m+n} D_i B_i^{m+n}(t), \tag{6}$$

where

$$D_i = \sum_{j=max(0,i-n)}^{min(m,i)} \frac{\binom{m}{j}\binom{n}{i-j}}{\binom{m+n}{i}} f_j g_{i-j}. \tag{7}$$

3. Polynomial approximation of rational curves

We need two-step to solve equation (2). One is for constrained control points of the approximation Bézier curve, the other is for unconstrained control points.

3.1. Constrained conditions of the approximation curve

Theorem 1. *Given a rational Bézier curve* $\mathbf{x}(t)$ *and a polynomial curve* $\tilde{\mathbf{y}}(t)$ *as in equations (1) and (2), if the following*

$$\begin{aligned}\mathbf{q}_r = \frac{1}{\omega_0\binom{m}{r}} \Bigg\{ &\binom{n}{r}\Delta^r\mathbf{P}_0 - \\ &\sum_{i=0}^{r-1}\left[\binom{m}{i}\binom{n}{r-i}\omega_{r-i}\mathbf{q}_i + (-1)^{r+i}\binom{m+n}{r}\binom{r}{i}\mathbf{C}_i^0\right]\Bigg\}\end{aligned} \tag{8}$$

$$\begin{aligned}\mathbf{q}_{m-s} = \frac{(-1)^s}{\omega_n\binom{m}{s}} \Bigg\{ &\binom{n}{s}\Delta^s\mathbf{P}_{n-s} - \\ &\sum_{i=1}^{s}\left[\binom{m}{s-i}\binom{n}{i}\omega_{n-i}\mathbf{q}_{m-s+i} - (-1)^{s+i}\binom{m+n}{s}\binom{s}{i}\mathbf{C}_{m+n-s+i}^1\right]\Bigg\}\end{aligned} \tag{9}$$

is true, where $0 < r + s < min(m, n)$, $\mathbf{C}_i^0 = \sum_{j=0}^{i} \frac{\binom{m}{j}\binom{n}{i-j}}{\binom{m+n}{i}}\omega_{i-j}\mathbf{q}_j$ *and* $\mathbf{C}_{m+n-s+i}^1 = \sum_{j=m-s+i}^{m} \frac{\binom{m}{j}\binom{n}{s+j-m-i}}{\binom{m+n}{s-i}}\omega_{m+n-s+i-j}\mathbf{q}_j$, *then we call* $\mathbf{x}(t)$ *and* $\tilde{\mathbf{y}}(t)$ *satisfy* $C^{(r,s)}$*-continuity.*

Proof. From equations (1), (6) and (7), we obtain

$$\begin{aligned}\sum_{i=0}^{n}\mathbf{P}_i B_i^n(t) &= \sum_{j=0}^{m}\mathbf{q}_j B_j^m(t)\sum_{i=0}^{n}\omega_i B_i^n(t) \\ &= \sum_{i=0}^{m+n}\mathbf{C}_i B_i^{m+n}(t),\end{aligned}$$

where

$$\mathbf{C}_i = \sum_{j=\max(0,i-n)}^{\min(m,i)} \frac{\binom{m}{j}\binom{n}{i-j}}{\binom{m+n}{i}}\omega_{i-j}\mathbf{q}_j. \tag{10}$$

Differentiating above equation and letting $t = 0$ yield

$$\binom{n}{r}\Delta^r \mathbf{P}_0 = \binom{m+n}{r}\Delta^r \mathbf{C}_0^0. \tag{11}$$

Here $\Delta^r \mathbf{C}_0^0$ can be rewritten as

$$\Delta^r \mathbf{C}_0^0 = (E - I)^r \, \mathbf{C}_0^0 = \sum_{i=0}^{r-1} (-1)^{r+i} \binom{r}{i} \mathbf{C}_i^0 + \mathbf{C}_r^0. \tag{12}$$

Since $0 < r + s < min(m, n)$, we can obtain the following result by equation (10)

$$\begin{aligned} \mathbf{C}_k^0 &= \sum_{j=0}^{k} \frac{\binom{m}{j}\binom{n}{r-j}}{\binom{m+n}{r}} \omega_{r-j} \mathbf{q}_j \\ &= \sum_{j=0}^{r-1} \frac{\binom{m}{j}\binom{n}{r-j}}{\binom{m+n}{r}} \omega_{r-j} \mathbf{q}_j + \frac{\binom{m}{r}}{\binom{m+n}{r}} \omega_0 \mathbf{q}_r. \end{aligned} \tag{13}$$

Finally, bring equations (12) and (13) into equation (11), which establish the formula (8).

Similarly, we can prove the formula (9) as $t = 1$. □

3.2. Unconstrained control points of the approximation curve

Differentiating equation (3) with respect to $\tilde{\mathbf{q}}_k$ yields the following equation:

$$\frac{\partial d}{\partial \tilde{\mathbf{q}}_k} = 2 \int_0^1 \rho(t) t^{r+1} (1-t)^{s+1} \left(\mathbf{x}(t) - \tilde{\mathbf{y}}(t) \right) J_k^{(r+1,s+1)}(2t-1) dt = \mathbf{0}.$$

Let

$$\rho(t) = \frac{\omega(t)}{t^{s+1}(1-t)^{r+1}},$$

we obtain

$$\int_0^1 (\mathbf{P}(t) - \omega(t)\tilde{\mathbf{y}}(t)) J_k^{(r+1,s+1)}(2t-1)dt = \mathbf{0}.$$

Substituting equation (2) into the above equation gives

$$\begin{aligned}
&\int_0^1 \mathbf{P}(t) J_k^{(r+1,s+1)}(2t-1)dt - \int_0^1 \omega(t) J_k^{(r+1,s+1)}(2t-1) \sum_{i=0}^{m} \mathbf{Q}_i B_i^m(t)dt \\
&= \int_0^1 t^{r+1}(1-t)^{s+1}\omega(t) J_k^{(r+1,s+1)}(2t-1) \sum_{j=0}^{m-(r+s+2)} \tilde{\mathbf{q}}_j J_j^{(r+1,s+1)}(2t-1)dt, \\
&k = 0, ..., m-(r+s+2)
\end{aligned} \tag{14}$$

where

$$\mathbf{Q}_i = \begin{cases} \mathbf{q}_i & i = 0, ..., r \text{ and } i = m-s, ..., m, \\ 0 & \text{others.} \end{cases}$$

For more detailed expressions, the first item of the equation (14) can be rewritten as using equations (4), (5) and (6)

$$\begin{aligned}
&\int_0^1 \mathbf{P}(t) J_k^{(r+1,s+1)}(2t-1)dt = \int_0^1 \sum_{i=0}^{n} \mathbf{P}_i B_i^n(t) \cdot \sum_{j=0}^{k} A_i B_i^k(t)dt \\
&= \frac{1}{n+k+1} \sum_{i=0}^{m+k} \left(\sum_{j=\max(0,i-n)}^{\min(k,i)} \frac{\binom{k}{j}\binom{n}{i-j}}{\binom{k+n}{i}} A_j \mathbf{P}_{i-j} \right), \\
&k = 0, .., m-(r+s+2).
\end{aligned}$$

Similarly, the second item of equation (14) can be rewritten as

$$\int_0^1 J_k^{(r+1,s+1)}(2t-1)\omega(t)\sum_{i=0}^{m}\mathbf{Q}_i B_i^m(t)dt$$
$$=\int_0^1\sum_{j=0}^{k}A_iB_i^k(t)\cdot\sum_{i=0}^{m+n}\tilde{\mathbf{C}}_iB_i^{m+n}(t)dt$$
$$=\frac{1}{m+n+k+1}\sum_{i=0}^{m+n+k}\left(\sum_{j=\max(0,i-m-n)}^{\min(k,i)}\frac{\binom{k}{j}\binom{m+n}{i-j}}{\binom{m+n+k}{i}}A_j\tilde{\mathbf{C}}_{i-j}\right),$$
$$k=0,..,m-(r+s+2),$$

where

$$\tilde{\mathbf{C}}_i=\sum_{j=\max(0,i-n)}^{\min(m,i)}\frac{\binom{m}{j}\binom{n}{i-j}}{\binom{m+n}{i}}\mathbf{Q}_j\omega_{i-j},$$

and for the left-hand side of equation (14), we have

$$\int_0^1 t^{s+1}(1-t)^{r+1}\sum_{i=0}^{n}\omega_iB_i^n(t)\sum_{i=0}^{k}A_iB_i^k(t)\sum_{j=0}^{m-(r+s+2)}\tilde{\mathbf{q}}_jJ_j^{(r+1,s+1)}(2t-1)dt$$
$$=\int_0^1 t^{s+1}(1-t)^{r+1}\sum_{j=0}^{m-(r+s+2)}\tilde{\mathbf{q}}_jJ_j^{(r+1,s+1)}(2t-1)\sum_{i=0}^{n+k}M_iB_i^{n+k}(t)dt$$
$$=\sum_{j=0}^{m-(r+s+2)}\sum_{i=0}^{n+k}M_i\tilde{\mathbf{q}}_j\int_0^1 t^{s+1}(1-t)^{r+1}J_j^{(r+1,s+1)}(2t-1)B_i^{n+k}(t)dt$$
$$=\sum_{j=0}^{m-(r+s+2)}\sum_{i=0}^{n+k}\sum_{l=0}^{j}\frac{(-1)^{j+l}\binom{j+r+1}{l}\binom{j+s+1}{j-l}\binom{n+k}{i}}{(n+k+r+s+j+3)\binom{n+k+r+s+j+2}{i+s+l+1}}M_i\tilde{\mathbf{q}}_j,$$
$$k=0,..,m-(r+s+2).$$

Since the Bernstein polynomials be linearly independent, the solution of linear system (14) is unique.

3.3. Further Approximation

When the weights $\omega_i = 1$ $(i = 0, ..., n)$ and the degree $m(< n)$ in equation (3), the polynomial approximation of rational Bézier curves degenerate into degree reduction of Bézier curves. We use this method to obtain smaller approximation error sometimes. That is, rational Bézier curves firstly approximated by Jacobi polynomial curves with a higher degree m', then we reduce the the Jacobi polynomial curves to degree m, where $n < m < m'$. The method can be summarized as the following algorithm:

Algorithm 1.

Step 1: Compute $\{\mathbf{q}\}_{i=0}^{r}$ and $\{\mathbf{q}\}_{i=m-s}^{m}$ using equation (8) and (9).

Step 2: Compute a Jacobi curve of degree m' using equation (14).

Step 3: Reduce the Jacobi curve to degree m using equation (14).

Step 4: Convert Jacobi curve to Bézier curve using equation (5), we obtain

$$\mathbf{y}(t) = \sum_{i=0}^{m} \mathbf{q}_i B_i^m(t), \tag{15}$$

where $\mathbf{q}_i$ were given by equations (5), (8) and (9).

3.4. Error estimation

Here, we present and prove a bound on the magnitude of the rational Bézier curve approximated by Bézier curve. A simple lemma was used as following [18].

Lemma 1. *If* $a_k \in \mathbb{R},\ b_k > 0$ *and* $c_k > 0,\ 1 \leq k \leq n$, *then*

$$\frac{\sum a_k c_k}{\sum b_k c_k} \leqslant \max_k \frac{a_k}{b_k}$$

Theorem 2. *For the rational Bézier curve and Bézier curve respectively*

defined by (1) and (15), we have

$$\|\mathbf{x}(t)-\mathbf{y}(t)\| \leqslant \max_i \frac{\sum\limits_{j=\max(0,i-m)}^{\min(n,i)} k_{ij}\,\|\mathbf{p}_j-\mathbf{q}_{i-j}\|}{\sum\limits_{j=\max(0,i-m)}^{\min(n,i)} k_{ij}} \leqslant \max_i \max_j \|\mathbf{p}_j-\mathbf{q}_{i-j}\|$$

where

$$k_{ij}=\binom{n}{j}\binom{m}{i-j}\omega_j.$$

Proof.

$$\begin{aligned}
&\|\mathbf{x}(t)-\mathbf{y}(t)\| \\
&= \left\| \frac{\sum\limits_{i=0}^{n}\omega_i\mathbf{p}_i B_i^n(t)-\sum\limits_{i=0}^{n}\omega_i B_i^n(t)\sum\limits_{j=0}^{m}\mathbf{q}_j B_j^m(t)}{\sum\limits_{i=0}^{n}\omega_i B_i^n(t)} \right\| \\
&\leqslant \frac{\sum\limits_{i=0}^{m+n}\sum\limits_{j=\max(0,i-m)}^{\min(n,i)}\frac{\binom{n}{j}\binom{m}{i-j}}{\binom{m+n}{i}}\omega_j\,\|\mathbf{p}_j-\mathbf{q}_{i-j}\|B_i^{m+n}(t)}{\sum\limits_{i=0}^{m+n}\sum\limits_{j=\max(0,i-m)}^{\min(n,i)}\frac{\binom{n}{j}\binom{m}{i-j}}{\binom{m+n}{i}}\omega_j B_i^{m+n}(t)} \\
&\leqslant \max_i \frac{\sum\limits_{j=\max(0,i-m)}^{\min(n,i)}\binom{n}{j}\binom{m}{i-j}\omega_j\,\|\mathbf{p}_j-\mathbf{q}_{i-j}\|}{\sum\limits_{j=\max(0,i-m)}^{\min(n,i)}\binom{n}{j}\binom{m}{i-j}\omega_j} \\
&\leqslant \max_i \max_j \|\mathbf{p}_j-\mathbf{q}_{i-j}\|,
\end{aligned}$$

which is completed the proof. □

4. Numerical examples

In this section, we provide two examples of application of the discussed methods. For each example, we give the Hausdoff distance.



Figure 1: The photomicrograph



Figure 2: The photomicrograph

Example 1. (Also Example 3 in [5, 7])The given curve is a rational Bézier curve of degree 9 with the control points $(17, 12)$, $(32, 34)$, $(-23, 24)$, $(33, 62)$, $(-23, 15)$, $(25, 3)$, $(30, -2)$, $(-5, -8)$, $(-5, 15)$, $(11, 8)$ and the associated weights $1, 2, 3, 6, 4, 5, 3, 4, 2, 1$. We find a 10-degree Bzier curve satisfying $C^{(0,0)}$-continuity with the given curve. The approximation errors under the Hausdorff distance from our method, Hu's method, Lewanowicz's method are 0.2424, 0.246726, and 0.317210, respectively. The resulting curves are shown in the Fig. 1 and the corresponding error distance curves are illustrated in the right-hand side of Fig. 2.

Example 2. (Also Example 3 in [7]) The given curve is a rational bézier curve of degree 8 with the control points $(0, 0)$, $(0, 2)$, $(2, 10)$, $(4, 6)$, $(6, 6)$, $(11, 16)$, $(8, 1)$, $(9, 1)$, $(10, 0)$ and the associated weights $1, 2, 3, 9, 12, 20, 30, 4, 1$. We find a 5-degree Bézier curve satisfying $C^{(1,1)}$-continuity with the given curve. The approximation errors under the Huasdorff distance from our method and Hu's method respectively are 0.4982 and 0.560612. The resulting curve of our method is shown in the Fig. 3 and the corresponding error distance curve is illustrated int the Fig. 4.


Original curve
C(0,0) with hybrid polynomials method
C(0,0) with degree reduction method
C(1,1) with hybrid polynomials method

Figure 3: The photomicrograph of Fig. 1



Figure 4: The photomicrograph of Fig. 1

Acknowledgements

This work is supported by the Natural Science Basic Research Plan in Shaanxi Province of China (No. 2013JM1004) and the Fundamental Research Funds for the Central Universities (No. GK201102025).

References

[1] L.A. Piegl, W. Tiller, K. Rajab, It is time to drop the "R" from NURBS, Engineering with Computers, 30 (4) (2014) 703-714.

[2] T. W. Sederberg, M. Kakimoto, Approximating rational curves using polynomial curves, in: G. Farin(Ed.), NURBS for curve and surface Design, SIAM, Philadelphia, 1991, 149-158.

[3] G. J. Wang, T. W. Sederberg, F. Chen, On the convergence of polynomial of rational functions, Journal Approximation Theory 89 (1997) 267-288.

[4] B. G. Lee, Y. B. Park, Approximate conversion of rational Bézier curves,

The journal of the Korean Society for Industrial and Applied Mathematics 1 (1998) 88-93.

[5] S. Lewanowicz, P. Woźny, P. Keller, Polynomial approximation of rational Bézier curves with constraints, Numerical Algorithms 59 (4) (2012) 607-622.

[6] M. Shi, J. S. Deng, Polynomial approximation of rational Bézier curves, Journal of Computational Information Systems 11 (20) (2015) 7339-7346.

[7] Q. Q. Hu, H. X. Xu, Constrained polynomial approximation of rational Bézier curves using reparameterization, Journal of Computational and Applied Mathematics 249 (2013) 133-143.

[8] A. R. Forrest, Interactive interpolation and approximation by Bzier curve, The Computer Journal 15 (1)(1972), 71-79.

[9] G. Frain, Curves and surfaces for CAGD, A Practical Guide, 5th edition, Morgan-Kaufmann, 2002.

[10] H. J. Kim, Y. J. Ahn, Good Degree Reduction of Bézier Curves Using Jacobi Polynomials, PERGAMON Computers and Mathematics with Applications 40 (2000), 1205-1215.

[11] G. Chen, G. J. Wang, Optimal multi-degree reduction of Bzier curves with constraints of endpoints continuity. Computer Aided Geometric Design 19 (2002) 365-377.

[12] Y. J. Ahn, Using Jacobi polynomials for degree reduction of Bzier curves with Ck-constraints, Computer Aided Geometric Design 20 (2003) 423-434

[13] J. M. Zheng, G. Z. Wang,Perturbing Bézier coefficients for best constrained degree reduction in the L_2-norm, Graphical Models 65 (2003) 351-368

[14] A. Rababah, Jacobi-Bernstein basis transformation, Comput. Meth. Appl. Math., 4 (2004) 206-214.

[15] A. H. Bhrawy, E.H. Doha, M.A. Saker, D. Baleanu, Modified JacobiCBernstein basis transformation and its application to multi-degree reduction of Bzier curves, Journal of Computational and Applied Mathematics 302 (2016) 369-384.

[16] G. Szego, Orthogonal polynomials, forth ed. American Mathemeatical Society, 1975.

[17] R. T. Farouki, V.T. Rajan, Algorithms for polynomials in Bernstein form, Computer Aided Geometric Design 5 (1988) 1-26.

[18] J. C. Kuang, Applied Inequalities 3rd, Sahngdong Science and technology Press, 2004, (in chinese).